

Treatment Guidelines for Major Depression in Older Adults



CCSMH
Canadian Coalition for
Seniors' Mental Health

CCSMPA
Coalition canadienne pour la
santé mentale des personnes âgées

Prevention

Assessment

Stepped-Care Treatment

For all older adults:

- Reducing isolation
- Physical activity
- Eating well
- Social activities
- Community involvement
- Mindfulness
- Reminiscence Therapy
- Self-Help CBT

In older adults consider:

- Prolonged bereavement
- Social isolation
- Chronic illness
- Memory difficulties
- Recurrent or prolonged hospitalization
- Dx of Parkinson's, dementia or stroke
- Recent placement in long term care
- Sleep, self-care and diet
- Recent onset anxiety

Psychosocial

For example:

- Reducing isolation
- Physical activity
- Eating well
- Mindfulness

Psychotherapy

For example:

- Brief Supportive counseling
- Cognitive Behaviour Therapy
- Problem Solving Therapy

Stepped Care Treatment focuses on using the least intensive and most effective level of intervention with regular reassessment.

Education/Self-Help

For example:

- Psychoeducation
- Self-Help books
- Online Self-Help CBT

Psychopharmacology

- Duloxetine or sertraline are recommended as first-line
- Escitalopram or citalopram are alternatives
- Consider anti-cholinergic effects and drug interactions

For complete treatment guidelines visit www.ccsmh.ca



Financial contribution from



Public Health
Agency of Canada

Agence de la santé
publique du Canada

This infographic is intended for educational purposes and not as a treatment plan. For the complete, updated treatment guidelines please scan the code or visit www.ccsmh.ca/projects/depression