

#### ccsmh.ca

# Canadian Guidelines for the Assessment and Treatment of Anxiety in Older Adults (2024)



### Recommended pharmaceutical interventions

Canadian Guidelines recommendations 28 and 30



Starting dose	Therapeutic dose	Maximum dose	Considerations					
First Line (Any of the following)								
2.5-5mg daily	10-20mg daily	10mg* daily	QTc Prolongation					
5-10mg daily	20-30mg daily	20mg* daily						
25-50mg daily	50-200mg daily	200mg daily	Indications in GAD, Panic, SAD					
37.5mg daily	150-300mg daily	300mg daily						
Second Line								
30mg daily	60-120mg daily	120mg daily	CYP2D6 inhibitor and substrate, risk for drug-drug interactions					
5mg BID to TID	10mg TID	10mg TID	In moderate anxiety, if first line not tolerated					
	First Line 2.5–5mg daily 5–10mg daily 25–50mg daily 37.5mg daily	First Line (Any of the following)  2.5–5mg daily  10–20mg daily  5–10mg daily  25–50mg daily  50–200mg daily  37.5mg daily  150–300mg daily  Second Line  30mg daily  60–120mg daily	First Line (Any of the following)  2.5-5mg daily  10-20mg daily  10mg* daily  5-10mg daily  20-30mg daily  20mg* daily  25-50mg daily  50-200mg daily  37.5mg daily  150-300mg daily  300mg daily  Second Line  30mg daily  120mg daily					

<sup>\*</sup>Health Canada maximum recommended dose based on evidence for QTc prolongation.

For additional medication information please visit GeriMedRisk



Financial contribution:





#### ccsmh.ca

## Canadian Guidelines for the Assessment and <u>Treatment of Anxiety in Older Adults (2024)</u>



### Pharmaceutical interventions not routinely recommended (except in specific circumstances)

Canadian Guidelines recommendations 29, 31, and 32

	Medication	Starting dose	Therapeutic dose	Maximum dose	Considerations
	Quetiapine fumarate extended release	50 mg daily	100-200 mg daily	300 mg daily	Poorly tolerated in frail older adults
	Quetiapine fumarate	12.5-25 mg once to twice daily	50-100 mg BID	150 mg BID	
	Pregabalin	25 mg daily	75-150 mg BID	150 mg BID	Tolerability issues, limited evidence for efficacy
	Lorazepam	0.25-0.5mg once daily	0.25-0.5mg BID	Not to exceed 2mg daily	Short-term, time-limited
	Clonazepam	0.125mg-0.25mg daily	O.125mg-O.25mg BID	Not to exceed 1mg daily	Long-acting, to be avoided in older adults

For additional medication information please visit GeriMedRisk



Financial contribution:

